

Fifth Int. Conference of Analysis and Applied Mathematics (ICAAM 2020)
23 September - 30 September 2020, Near East University, Turkey

On the boundedness of solution of the parabolic differential equation with time involution

Allaberen Ashyralyev^{1,2,3}, Amer Ahmed¹

¹Department of Mathematics, Near East University, Nicosia, TRNC, Mersin 10, Turkey

²Peoples Friendship University Russia, Ul Miklukho Maklaya 6, Moscow 117198, Russia

³Institute of Mathematics and Mathematical Modeling, 050010, Almaty, Kazakhstan,

aallaberen@gmail.com

amermzory81@gmail.com

In the present paper, the initial value problem for the parabolic type involutory partial differential equation is investigated. Applying Green's function of space operator generated by the differential problem, we get formula for solution of this problem. In applications, the theorem on stability estimates for the solution of this problem is established. Furthermore, it is applied to the existence and uniqueness of bounded solution of the nonlinear problem with involution.

KEYWORDS: Hilbert space, Parabolic equation with involution; Boundedness.

2010 Mathematics Subject Classification: 35J25, 47E05, 34B27.

References

- [1] A.Sarsenbi, Well-Posedness of Mixed Problems for Parabolic type with Involution, PhD Thesis, CSU, Kazakhstan, 2019.
- [2] A. Ashyralyev, T. Abbas, A numerical algorithm for the involutory parabolic problem, Third International Conference of Mathematical Sciences (ICMS 2019) AIP Conf. Proc. 2183, 070015-1–070015-3; <https://doi.org/10.1063/1.5136177>.
- [3] A.Cabada, F. Tojo, Differential Equations with Involutions. Atlantis Press, 2015.
- [4] A. Ashyralyev, P.E. Sobolevskii, New Difference Schemes for Partial Differential Equations, Operator Theory Advances and Applications, Birkhauser Verlag, Basel, Boston, Berlin, 2004.