

Fourier method approach in mixed problems for the heat equation with involution perturbation

Sarsenbi Abdisalam ¹

¹ *M. Auezov South Kazakhstan State University, Shymkent, Kazakhstan
abdisalam@mail.ru*

Abstract: In this paper we consider a mixed problem for the parabolic equation with involution of the form

$$u_t(x, t) = u_{xx}(x, t) - \alpha u_{xx}(-x, t) - q(x)u(x, t), \quad -1 < x < 1, \quad t > 0; \quad (1)$$

$$u(0, x) = \varphi(x), \quad -1 \leq x \leq 1; \quad u(-1, t) = 0, \quad u(1, t) = 0, \quad t \geq 0. \quad (2)$$

In order to solve the problem we apply the separation of variables supposing $u(x, t)$ is a product:

$$u(x, t) = X(x)T(t).$$

Here the function $X(x)$ gives the solution to the boundary value problem

$$-X''(x) + \alpha X''(-x) + q(x)X(x) = \lambda X(x), \quad -1 < x < 1, \quad |\alpha| < 1,$$

$$X(-1) = 0, \quad X(1) = 0. \quad (3)$$

Theorem 1. If the number $\sqrt{\frac{1-\alpha}{1+\alpha}}$ is not even then the system of eigenfunctions of the spectral problem (3) forms the basis in $L_2(-1, 1)$.

Theorem 2. Let the following conditions be satisfied:

- 1) a real continuous function $q(x)$ is non-negative;

2) the number $\sqrt{\frac{1-\alpha}{1+\alpha}}$ is not even ($-1 < \alpha < 1$);

3) the initial condition $\varphi(x)$ of the mixed problem (1), (2) satisfies the conditions $\varphi(-1) = \varphi(1) = 0$,

Then the solution to the mixed problem (1), (2) exists, is unique and can be represented by the series

$$u(x, t) = \sum_{k=1}^{\infty} A_k e^{-\lambda_k t} X_k(x).$$

1. ACKNOWLEDGEMENT

This work was supported by the Science Committee of the Ministry of Education and Science of the Republic of Kazakhstan, grant AP05131225.

REFERENCES

- [1] A.M. Sarsenbi; *The theorem on the basis property of eigenfunctions of second order differential operators with involution*, AIP Conference Proceedings, vol. 1759, p. 020030, doi: 10.1063/1.4959644, 2016.
- [2] L.V. Kritskov, A.M. Sarsenbi; *Riesz basis property of system of root functions of second-order differential operator with involution*, Differential Equations, vol. 53. no. 1. pp. 33-46, 2017.